watsons






# Introduction

Zika virus (ZIKV) infection in humans has been reported since the 1950s. Zika virus infection is a **mosquito-borne infection** and has **symptoms similar to Dengue**. Singapore's first imported case of Zika virus infection was confirmed on 13<sup>th</sup> May 2016, and we should exercise vigilance to prevent the spread of the Zika virus infection.

## Transmission

- Zika virus is transmitted via the *Aedes* mosquito, the same species that is responsible for the spread of Dengue virus.
- 4 main modes of transmission:
  - Via the bite of an infected Aedes mosquito
  - > Maternal-fetal (either through the placenta or during delivery)
  - Sexual transmission




#### Diagnosis

- To date, the Zika virus infection can only be confirmed through a blood test.
- See your doctor if you develop any of the signs and symptoms mentioned below and have visited an area where Zika is found.
- If you have recently travelled, inform your doctor when and where you have travelled.

### Signs and symptoms

- Only 1 in 5 infections are symptomatic.
- Zika virus infection is generally a mild and self-limiting illness.
- Symptoms of Zika are similar to those of **Dengue**: .



### Areas with Zika

Refer to the Ministry of Health (MOH) website for an updated list of areas with Zika virus outbreak / ongoing local transmission

https://www.moh.gov.sg/content/moh\_web/home/pressR oom/Current Issues/2016/zika-virus.html



- There is no medicine to specifically treat the Zika virus infection.
- Treat the symptoms:
  - Get plenty of rest  $\checkmark$
  - Prevent dehydration by drinking enough water  $\checkmark$
  - Paracetamol may be useful in reducing fever or  $\checkmark$ pain
  - Do not take aspirin or other non-steroidal anti-1 inflammatory drugs (NSAIDS) until dengue can be ruled out to reduce the risk of bleeding


### Complications

The Zika virus infection has been known to result in microcephaly (where baby's head is born smaller than normal) and Guillain-Barré syndrome (a disorder of the nervous system).

### Prevention

- There is currently no vaccine to prevent the Zika virus infection.
- 3 main ways to prevent the spread of the Zika virus:

#### 1. Remove stagnant water



- 2. Avoid mosquito bites by applying insect repellents / wearing adequate clothing for protection.
- 3. Postpone travel to areas with ongoing outbreak or local transmission

### Zika & Pregnancy

- Zika virus infection during pregnancy has been linked to babies born with microcephaly (where baby's head is much smaller than normal).
- Zika virus infection in a woman who is NOT pregnant will not pose a risk for birth defects in future pregnancies after the virus has cleared from her blood.



Baby with Microcephaly

Prevent sexual transmission of the Zika virus by ensuring that condoms are used correctly from start to finish, every time during sex.

Updated on 3<sup>rd</sup> June 2016

#### Disclaimer

Information provided by this patient information leaflet is solely for informational purposes only and is not intended as a substitute for the advice provided by your physician, pharmacist or other healthcare professional. You should not use the information for diagnosis or treatment of a health problem or disease. Always speak with your physician, pharmacist or other healthcare professional before taking any medicine or supplement, or adopting any treatment for a health problem. Under no circumstances will Watson's Personal Care Stores Pte Ltd be liable to any person for damages of any nature arising in any way from the use of such information.

Prepared by: Clara Lin, Pharmacist @ Watsons Changi Airport Terminal 3 Transit North

Vetted by: Chung Wing Lam, Principal Clinical Pharmacist @ Watsons Ngee Ann City

#### Images from:

- Baby with microcephaly -1.
- http://www.cdc.gov/ncbddd/birthdefects/microcephaly.html 2. 5-step mozzie wipeout - http://www.motherinc.org/seriously-
- somewhat-serious/the-mozzie-wipeout/

