

What is this medicine for?

Bupropion is an antidepressant used to treat symptoms of low mood such as sad feelings, feelings of hopelessness or worthlessness, feeling bad about yourself, suicidal thoughts, poor sleep, energy or appetite.

Bupropion is also used as an aid in quitting smoking.

The medicine may also be used for other conditions, check with your doctor.

Before using this medicine:

Tell your doctor if you have any drug allergies or have ever had heart, liver, or kidney disease; seizures (fits); eating disorders; diabetes; urinary problems; high cholesterol; low blood pressure; phenylketonuria; bipolar disorder; schizophrenia.

FOR WOMEN: Tell your doctor if you are pregnant, plan to become pregnant, or are breast-feeding.

How to use this medicine?

Follow the directions on your medicine label and take the medicine exactly as directed. Do not take more or less of it or take it more often than prescribed by your doctor. Bupropion can be taken with or without food.

Do NOT break, split or crush tablets. Swallow whole.

This medicine must be taken regularly for a few weeks before its full effect is felt. Continue to take this medicine regularly as prescribed even if you feel well. Do not stop taking it without talking to your doctor.

Once these symptoms are under control, you should continue taking this medication regularly to prevent the return of your depressive or anxiety

symptoms. Do check with your doctor on the appropriate length of treatment you need.

Stopping this medication suddenly after taking for a long time may lead to mild discontinuation symptoms such as nausea, anxiety, insomnia, headache and dizziness.

Precautions while taking this medicine:

- Bupropion may make you drowsy or less alert. Do not drive or operate machinery if affected.
- Avoid drinking alcohol while taking this medicine. Drinking alcohol can worsen symptoms of depression and side effect of drowsiness.
- When you take this medicine with some other medicines, it may affect how they work. Be sure that you tell all doctors, pharmacists, dentists and other health care providers who treat you about all medicines and natural remedies you are taking.
- Your mental condition may change in unexpected ways especially at the beginning (first few weeks) of treatment and during dose changes. Contact your doctor immediately if you or your child (the patient) experience new or worsening thoughts of harming yourself, increase in symptoms such as anxiety, restlessness, violent behaviour, or other changes in mood and behaviour. Be sure that your family or caregiver knows which symptoms may be serious so they can call the doctor if you are unable to seek treatment on your own.

What should I do if I forget a dose?

Take the missed dose as soon as you remember it. However, if it is more than half way between the doses, skip the missed dose and take your medicine at the usual time.

Do not take two or more doses at the same time to make up for a missed one.

Possible Side Effects

Like all medications, this medication may also cause some POSSIBLE side effects in some people. Everyone responds differently to medications; so not everyone experiences any or all of them.

COMMON (Generally affect up to 10% of patients)

Most of these side effects below will get better or go away completely with continued medication taking as your body gets used to this medication. Contact your doctor or pharmacist should they become too bothersome to you.

- ~ Headache
- ~ Nausea
- ~ Insomnia
- ~ Dry mouth (suck ice chips or sugarless candy if necessary)
- ~ Constipation (drink more fluids, regular exercise, take fiber-rich diet, or talk to your doctor about stool softener or laxative)
- ~ Dizziness (can be reduced by getting up slowly from a sitting or lying down position)
- ~ Tiredness

FOR URGENT ATTENTION

Check with your doctor as soon as possible if you have any of the following:

- ~ Allergic reaction: Skin rash or itching, difficulty breathing, swelling of the eyes, face, or lips
- ~ Swelling in hands, ankles or feet
- ~ Flu-like symptoms (fever, chills, sore throat); mouth sores / ulcers
- ~ Severe vomiting or diarrhoea, loss of appetite, abdominal pain
- ~ Confusion, severe weakness, uncontrolled movements or twitching of muscles
- ~ Feelings of intense agitation or anxiety, restlessness or inability to sit still, or irritability
- ~ Unusual thoughts; talking, feeling, and acting with excitement that you cannot control
- ~ Seizures (fits)
- ~ Severe or continuing headache, or ringing in the ears

If you have any concerns about your medications or if you have other side effects that you think are caused by this medicine, please consult with your doctor or pharmacist.

Where should I store the medicine?

Store this medicine at room temperature, away from direct light, heat and moisture (not in the bathroom) and out of reach of children.

Throw away any medication that is expired or no longer needed.

Do not remove the tablet from the blister until just prior to taking your tablet; once removed, it cannot be stored.

Do not share the medicine with other people.

Ask your doctor or pharmacist for more information or if you have any questions.

Institute of Mental Health

Pharmacy Department

10 Buangkok View
Singapore 539747

For medication enquiries: (during office hours)

IMH Clinic B/ Sayang Wellness Pharmacy: 63892077

IMH Clinic C (NAMS) Pharmacy: 63893666

Community Wellness Centre Pharmacy: 64715797

Geylang Behavioural Medicine Pharmacy: 67461724

Child Guidance Clinic (HPB) Pharmacy: 64353807

Child Guidance (Sunrise Wing) Pharmacy: 63892410

Email: enquiry@imh.com.sg



Patient Information Leaflet on

BUPROPION