

What is this medicine for?

Paliperidone is used to treat and prevent return of psychosis (such as hearing or seeing things which are actually not present, confused or unreal thoughts or uncontrolled anger).

Paliperidone may also be used for other conditions, check with your doctor.

Before using this medicine:

Tell your doctor if you have or have ever had Parkinson's disease; Alzheimer's disease; Dementia; heart or kidney disease; stroke; diabetes; severe pre-existing gastrointestinal narrowing or other disorders; difficulty swallowing; seizures (fits); breast cancer or drug allergies.

FOR WOMEN: Tell your doctor if you are pregnant, plan to become pregnant, or are breast-feeding.

Precautions while taking this medicine:

- Paliperidone may make you drowsy or less alert. Do not drive or operate machinery if affected. Avoid alcohol.
- Paliperidone may sometimes worsen blood glucose control in diabetes. Maintain regular check-ups with your doctor.
- Paliperidone may interfere with your body's ability to cool down (make you sweat less), causing your body temperature to increase. Use extra care not to become overheated during strenuous exercise or hot weather.

- When you take this medicine with some other medicines, it may affect how they work. Be sure that you tell all doctors, pharmacists, dentists and other health care providers who treat you about all medicines and natural remedies you are taking.

How to use this medicine?

Follow the directions on your medicine label and take the medicine exactly as directed. Do not take more or less of it or take it more often than prescribed by your doctor.

The extended-release tablets are to be swallowed whole with water or other liquids. Do not break, cut, crush or chew the tablet before swallowing.

It is normal to notice the tablet shell in the stools sometimes. This is because the empty tablet shell left after the medicine has been absorbed into your body does not dissolve completely.

Continue to take this medicine as prescribed even if you feel well. Do not stop taking it without talking to your doctor, otherwise your condition may worsen very quickly. Paliperidone must be taken regularly for a few weeks before its full effect is felt.

For patients on Paliperidone (Invega Sustenna™) injection:

- It is given as an injection into your upper arm or buttock muscle.

What should I do if I forget a dose?

Take the missed dose as soon as you remember it. However, if it is more than half way between the doses, skip the missed dose and take your medicine at the usual time.

Do not take two doses at the same time to make up for a missed one.

For patients on Paliperidone (Invega Sustenna™) injection:

- Please call up your doctor to arrange for an appointment visit.

Possible Side Effects

Like all medications, this medication may also cause some POSSIBLE side effects in some people. Everyone responds differently to medications; so not everyone experiences any or all of them.

COMMON (Generally affect up to 10% of patients)

Most of these side effects below will get better or go away completely with continued medication taking as your body gets used to this medication. Contact your doctor or pharmacist should they become too bothersome to you.

- ~ Headache
- ~ Nausea / Vomiting
- ~ Increased salivation
- ~ Drowsiness
- ~ Dizziness (can be reduced by getting up slowly from a sitting or lying down position)
- ~ Weight gain
- ~ Mild pain, itchiness, redness or swelling at the injection site (For Paliperidone injection only)

FOR URGENT ATTENTION

Check with your doctor as soon as possible if you have any of the following:

- ~ Allergic reaction: Skin rash or itching, difficulty in breathing, swelling in the face, lips or other parts of the body, or chest tightness
- ~ Fast or irregular heartbeat
- ~ High fever, muscle stiffness, sweating or confusion
- ~ Difficulty in speaking or swallowing
- ~ Muscle spasms (such as face, neck, back and eyelids), twitching or uncontrolled body movements
- ~ Restlessness or need to keep moving constantly
- ~ Seizures (fits)
- ~ Blurred vision

If you have any concerns about your medications or if you have other side effects that you think are caused by this medicine, please consult with your doctor or pharmacist.

Where should I store the medicine?

Store this medicine at room temperature, away from direct light, heat and moisture (not in the bathroom) and out of reach of children.

Throw away any medication that is expired or no longer needed.

For patients on Paliperidone (Invega Sustenna™) injection, this medication will be administered to you at a hospital or at your doctor's clinic. You should not store it at home.

Do not share the medicine with other people.

Ask your doctor or pharmacist for more information or if you have any questions.

Institute of Mental Health

Pharmacy Department

10 Buangkok View

Singapore 539747

For medication enquiries: (during office hours)

IMH Clinic B \ Sayang Wellness Pharmacy: 63892077

IMH Clinic C (NAMS) Pharmacy: 63893666

Community Wellness Centre Pharmacy: 64715797

Geylang Behavioural Medicine Pharmacy: 67461724

Child Guidance Clinic (HPB) Pharmacy: 64353807

Child Guidance (Sunrise Wing) Pharmacy: 63892410

Email: enquiry@imh.com.sg



Patient Information Leaflet on

PALIPERIDONE

